

# The Thirtieth Annual Eastern Shore High School Mathematics Competition

November 7, 2013

## Team Contest Exam

### Instructions

Answer as many questions as possible in the time provided. To receive full credit for a correct solution, show all work and provide a clearly written explanation. Solutions will be judged based on correctness, completeness and clarity. (Little credit, if any, will be given for a solution consisting of just a number or a single sentence.)

All work and answers must be written on the provided sheets of plain white paper. Use only one side of each sheet of paper, and start each new problem on a new sheet of paper. Write your team name (that is, the name of the school which you are representing) at the top of each sheet that you turn in for scoring.

**At the start of the team round, your team will receive a copy of only Problem 1. Your team must submit a response to Problem 1 within the first 15 minutes of the team round time interval.**

**When you submit your response for Problem 1, you will receive a copy of Problem 2 and a copy of Problem 3. Your team will then have the time remaining in the team round to complete a response for each problem.**

Note: if your team completes Problem 1 before the end of the allotted time, you may submit it and receive copies of Problem 2 and Problem 3 in advance.



2. Note that the set of **positive integers** is  $\{1; 2; 3; 4; 5; \dots; g\}$ .

Although  $\sqrt{2} + \sqrt{3}$  does not equal the square root of a positive integer,  $\sqrt{27} + \sqrt{48}$  does.

(a) Find a positive integer  $n$  such that  $\sqrt{27} + \sqrt{48} = \sqrt{n}$ .  
**You must provide written work to show why  $\sqrt{27} + \sqrt{48} = \sqrt{n}$ .**

(b) Find another example like the one in part (a).  
In other words, find positive integers  $a$ ,  $b$ , and  $c$  such that  $\sqrt{a} + \sqrt{b} = \sqrt{c}$ .  $a$ ,  $b$ , and  $c$  cannot be square numbers.  
**You must provide written work to justify your answer.**

(c) Find every set of positive integers  $a$ ,  $b$ , and  $c$  such that

$a$ ,  $b$ , and  $c$  are all less than 35

$$\sqrt{a} + \sqrt{b} = \sqrt{c}$$


$a$ ,  $b$ , and  $c$  are distinct

None of  $\sqrt{a}$ ,  $\sqrt{b}$ , and  $\sqrt{c}$  is an integer.

3. For two points,  $P = (x_1; y_1)$  and  $Q = (x_2; y_2)$  in a coordinate plane, the **Taxicab Distance** between  $P$  and  $Q$  is

$$d(P; Q) = |x_2 - x_1| + |y_2 - y_1|$$

Illustrated below, it measures the distance as a taxicab would travel on a rectangular grid of city streets.



(a) Find  $d(R; S)$  if  $R = (1; 6)$  and  $S = (4; 2)$ .

A **Taxicab Loop** is the set of all points that have the same Taxicab Distance from a given point, referred to as the **Dispatch Point**. The Taxicab Distance from the Dispatch Point to any point on the Loop is known as the **Radius of the Taxicab Loop**.

(b) Sketch a Taxicab Loop with Dispatch Point at  $(0, 0)$  and with Radius 1.

The **Taxicab Length** of a path consisting of straight line segments is the sum of the Taxicab Lengths of the segments. (a) The Taxicab Length of a path consisting of straight line segments is the sum of the Taxicab Lengths of the segments.